

Microstructural abnormalities of the brain white matter in attention-deficit/hyperactivity disorder

Lizhou Chen, PhD*; Xiaoqi Huang, MD, PhD*; Du Lei, PhD; Ning He, MD; Xinyu Hu, MM; Ying Chen, PhD; Yuanyuan Li, MD; Jinbo Zhou, MM; Lanting Guo, MD; Graham J. Kemp, MA, DM; Qiyong Gong, MD, PhD

Background: Attention-deficit/hyperactivity disorder (ADHD) is an early-onset neurodevelopmental disorder with multiple behavioural problems and executive dysfunctions for which neuroimaging studies have reported a variety of abnormalities, with inconsistencies partly owing to confounding by medication and concurrent psychiatric disease. We aimed to investigate the microstructural abnormalities of white matter in unmedicated children and adolescents with pure ADHD and to explore the association between these abnormalities and behavioural symptoms and executive functions. **Methods:** We assessed children and adolescents with ADHD and healthy controls using psychiatric interviews. Behavioural problems were rated using the revised Conners' Parent Rating Scale, and executive functions were measured using the Stroop Colour-Word Test and the Wisconsin Card Sorting test. We acquired diffusion tensor imaging data using a 3 T MRI system, and we compared diffusion parameters, including fractional anisotropy (FA) and mean, axial and radial diffusivities, between the 2 groups. **Results:** Thirty-three children and adolescents with ADHD and 35 healthy controls were included in our study. In patients compared with controls, FA was increased in the left posterior cingulum bundle as a result of both increased axial diffusivity and decreased radial diffusivity. In addition, the averaged FA of the cluster in this region correlated with behavioural measures as well as executive function in patients with ADHD. **Limitations:** This study was limited by its cross-sectional design and small sample size. The cluster size of the significant result was small. **Conclusion:** Our findings suggest that white matter abnormalities within the limbic network could be part of the neural underpinning of behavioural problems and executive dysfunction in patients with ADHD.

Introduction

Attention-deficit/hyperactivity disorder (ADHD) is one of the most common neurodevelopmental disorders in childhood; it has a reported prevalence of 5.3% in school-age children¹ and 2.5% in adults² worldwide. As well as behavioural problems (inappropriate hyperactivity and impulsivity, inattention symptoms), patients show deficits in executive functions (interference control, cognitive flexibility, working memory, planning and organization)³ that can impair academic and vocational performance.⁴ This disorder is a childhood-onset condition, so the pathophysiological profile in adult patients is more likely to be complicated by interactions with other factors during neuromaturation.⁵ Studies of children and adolescents with ADHD may therefore give a clearer view of its pathogenesis.

Neuroimaging methods have been useful in exploring ADHD pathophysiology. Much has been learned from macrostructural MRI about abnormalities in grey matter structures, such as the basal ganglia, prefrontal lobe, temporal and parietal cortices,⁶ and from fMRI about abnormal activation patterns in frontostriatal, frontotemporal and frontoparietal regions.⁷ This functional work has led to the notion of abnormal large-scale brain circuits in ADHD — notably the frontal–striatal–cerebellar circuit,⁸ with more recent evidence implicating the default mode (DMN), limbic and visual networks.⁹ Attention has also turned to white matter microstructural alterations in patients with ADHD⁸ and to models involving abnormal connections between multiple functional units in neural networks conceived as ensembles of neuronal bodies and axonal pathways.¹⁰ The

Correspondence to: Q. Gong, Huaxi MR Research Center, Department of Radiology, West China Hospital of Sichuan University, Chengdu 610041, China; qiyonggong@hmrc.org.cn

*These authors contributed equally to this work.

Submitted Jul. 18, 2014; Revised Nov. 15, 2014; Accepted Dec. 2, 2014; Early-released Apr. 8, 2015.

DOI: 10.1503/jpn.140199

©2015 8872147 Canada Inc.

importance of white matter deficits in patients with ADHD is supported by the observation that the effect sizes for volume reductions in total brain and lobar volumes are larger in white matter (0.30–0.64) than in grey matter (0.27–0.35).¹¹

Diffusion tensor imaging (DTI) offers a sensitive and neurobiologically relevant way to characterize microstructural white matter alterations *in vivo*.¹² Of several DTI-derived indices, fractional anisotropy (FA) is the most commonly used, reporting the local and regional directional coherence of water diffusion.¹³ Additional information is provided by the decomposition of FA into axial diffusivity (AD) and radial diffusivity (RD), which reflect diffusion parallel to and perpendicular to the axons, respectively, and by mean diffusivity (MD), which reflects the magnitude of diffusion in white matter pathways.¹³ Of the various analytic approaches to DTI, whole-brain analysis provides the best unbiased measure of brain regions that may be missed by region of interest (ROI) analysis based on a priori assumptions and thus offers greater reliability in screening distributed anomalies in patients with ADHD.¹⁴

Nine published DTI studies have used whole-brain approaches in children and adolescents with ADHD,^{5,15–22} reporting altered FA over diverse white matter structures. Such discrepancies may be owing to the differences in sample sizes, demographic characteristics and analysis pipelines (see the Appendix, Table S1, available at [jpn.ca](#)). Two of these factors deserve particular attention. Most of these studies included patients who either were currently taking medication or who had a history of taking medication; as medication has been reported to normalize white matter deficits,¹¹ the potential effect of medication is an important confounding factor. The other main confounding factor is comorbid presentation with other psychiatric and/or developmental disorders that may have unique etiological pathways or share pathophysiology with ADHD.²³

Another important issue about ADHD is the poorly understood association between the atypical brain circuitry revealed with neuroimaging methods and the executive dysfunctions measured with neuropsychological tests.²⁴ Although fMRI studies have suggested an association with the aberrant frontoparietal circuit,⁷ debate remains about which core deficits of brain function lead to the impairments of ADHD.²⁵ Diffusion tensor imaging may provide evidence of microstructural abnormalities in the brain circuits underlying specific executive deficits.

We therefore set out to study a relatively large sample of medication-naïve children and adolescents with pure ADHD, defining white matter microstructural abnormalities through a whole-brain DTI approach and testing for their correlations with clinical symptoms and executive dysfunctions.

Methods

Participants

We recruited patients with ADHD from the Department of Psychiatry at West China Hospital, Sichuan University. Diagnosis of ADHD was determined by 2 experienced clinical

psychiatrists (L.G. and N.H.; 28 and 5 yr of experience in clinical psychiatry, respectively) using the Chinese modified version of Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version (SCID-I, patient edition).²⁶ We excluded patients with oppositional defiant disorder, conduct disorder, Tourette disorder or any other Axis I psychiatric comorbid disorders. Other exclusion criteria were a full-scale IQ lower than 90 based on an age-appropriate Wechsler Intelligence Scale for Children — Chinese Revision,²⁷ current use of or history of taking psychotropic medication, left-handedness as assessed using the Annett Hand Preference Questionnaire,²⁸ pregnancy or substantial physical illness and standard MRI scanning contraindications.

We recruited healthy controls via advertisement in local schools; controls were also screened using the Chinese modified version of SCID-I (nonpatient edition) to exclude any Axis I psychiatric diagnosis and use of any psychotropic medications. Controls were required to have no family history of psychiatric illness in first-degree relatives. Other exclusion criteria were the same as those for the ADHD group.

Two neuroradiologists (L.C. and X.H., 3.5 and 3 yr of experience in neuroimaging, respectively) inspected conventional MRIs of all participants to exclude gross abnormalities.

Approval for this study was granted by the ethical committee of Sichuan University and the research ethics board at the West China Hospital of Sichuan University. All participants and their parents were fully informed about the purpose and procedures of this study, and written informed consent was obtained from the parents.

Behavioural measures and executive function tests

We used the Chinese version of the revised Conners' Parent Rating Scale (CPRS)²⁹ to measure the behavioural problems of ADHD patients. The CPRS is a battery of questions to evaluate problematic behaviour across areas such as sleep, temper and peer relationships. The parent rates each behaviour on a 4-point Likert-type scale; the behaviours are aggregated into 6 factors: conduct problems, study problems, psychosomatic, impulsive-hyperactive, anxiety and hyperactivity index. Higher scores indicate more severe problems.

The Stroop Colour-Word (Stroop-CW) Test³⁰ and the Wisconsin Card Sorting Test (WCST)³¹ are commonly used neuropsychiatric tests of executive abilities. The Stroop-CW Test, in which the participant is shown a list of names of colors printed in mismatching colors and is asked to name the colour instead of the word, is often used to measure executive interference control.³² The dependent variables are number right, number error, number correction and total time. The WCST (64-card version), in which the participant is asked to change matching strategies after correctly matching a card according to a certain stimulus feature (number, colour, form) for 10 consecutive trials, mainly reflects cognitive flexibility,³³ involving perceptual learning, set-shifting, working memory and executive control; the dependent variables are perseverative errors, nonperseverative errors, total errors and categories completed.

Data acquisition

Whole brain MRI data were acquired in the West China Hospital, Sichuan University, using a Siemens Trio 3 T MRI system with an 8-channel phase-array head coil. Sequences included a spoiled gradient-recalled echo sequence to acquire the T_1 -weighted images (repetition time [TR] 1900 ms, echo time [TE] 2.26 ms, section thickness 1 mm; matrix 256×256 ; field of view [FOV] 256 mm) and a single-shot echo-planar sequence (TR 6800 ms, TE 93 ms, section thickness 3 mm with no intersection gaps, matrix 128×128 , FOV 230 mm) to obtain DTIs in 50 axial planes with 20 noncollinear diffusion sensitization gradients ($b = 1000 \text{ s/mm}^2$) as well as a reference image with no diffusion weighting (b_0 image). Earplugs were used, and participants lay supine with their heads fixed by foam pads to minimize head movement. We took several steps to ensure optimal scan quality: good communication with participants and their parents allayed any worries and helped the participants cooperate during scanning. If monitoring detected any movement, the scanning sequence was halted and repeated; the acquired images were checked immediately by 2 neuroradiologists (L.C. and X.H.), and those with suspected artifacts were rescanned. Finally, during preprocessing all DTI maps were checked for artifacts in order to discard all data from the affected participants.

Statistical analysis

Diffusion-weighted images of each participant were analyzed using the Diffusion Toolbox FSL software version 1.1 (FMRIB Software Library). First, the data were corrected for head motion and eddy currents using affine registration to the b_0 image volume. Next, brain extraction was performed to delete non-brain tissue from the whole head image. Finally, we calculated parameter maps for FA, MD, AD and RD using "dtifit," which fits a diffusion tensor model to each voxel and estimates the principal directions of diffusion. Voxel-based analysis was then performed using SPM8 (Wellcome Trust Centre for Neuroimaging; <http://fil.ion.ucl.ac.uk>) running in MatLab 2012 (MathWorks). Each b_0 image was nonlinearly normalized using the echo-planar imaging template supplied with SPM8 to estimate the normalization parameter, which was applied to all parameter maps, each voxel being $2 \times 2 \times 2 \text{ mm}^3$. Finally, the normalized parameter maps were smoothed using an isotropic Gaussian filter (6 mm full-width at half-maximum).

We performed a voxel-wise comparison of FA and MD between patients with ADHD and controls using a 2-sample t test in SPM8 with age and sex as covariates. The threshold was taken as $p < 0.05$ for false discovery rate (FDR) correction. The resultant clusters were defined as ROIs through the MarsBaR toolbox in SPM8, and we extracted their averaged FA and MD values. We performed a 2-tailed Pearson correlation analysis in the ADHD group between the extracted values and multiple measurements, including CPRS, Stroop-CW Test and WCST results, using SPSS version 20.0 (IBM).

We extracted AD and RD values from clusters identified with significant FA differences through the MarsBaR toolbox and compared them between the ADHD and control groups using a 2 sample t test in SPSS.

As adolescents are undergoing rapid brain development, we performed a subgroup analysis on participants younger than 12 years. To minimize sex influences, we also performed a subgroup analysis on male participants.

Results

Participants

Our study included 33 patients aged 7–16 years with ADHD (all combined subtype) who were drug-naive and did not have any comorbidities. We also included 35 healthy controls aged 7–14 years. All participants were from the Chinese Han population and were right-handed. As shown in Table 1, demographic variables, including age, sex, IQ scores, birth weight and parental education years were not significantly different between the 2 groups.

Behavioural measures and executive function tests

Table 1 shows the results of behavioural measures and executive function tests. As expected, patients with ADHD had more severe behavioural problems and worse executive function than controls. In the CPRS, patients scored higher than controls in all indices ($p < 0.001$) except anxiety ($p = 0.30$). In the Stroop-CW test, patients had fewer right answers ($p < 0.001$), more errors ($p < 0.001$) and corrections ($p < 0.001$) and longer total time ($p < 0.001$) than controls. In the WCST, patients achieved fewer total correct responses ($p = 0.002$) and categories completed ($p = 0.001$) and made more total errors ($p = 0.001$), perseverative errors ($p = 0.001$) and nonperseverative errors ($p = 0.006$) than controls.

Group differences in diffusion characteristics and correlation analysis

As shown in Figure 1, relative to controls, patients with ADHD had 1 cluster showing increased FA in the left posterior cingulum bundle (MNI coordinates: $x, y, z = -6, -20, 32$, $t = 5.23$, cluster size 8 voxels). In no instance did patients with ADHD show significantly lower FA than controls, and there were no significant differences in MD between the ADHD and control groups. For the cluster with significant FA differences, the ADHD group had significantly higher AD and lower RD than controls (AD: ADHD 12.00 ± 0.60 v. control 11.42 ± 0.57 , $p < 0.001$; RD: ADHD 7.17 ± 0.53 v. control 7.42 ± 0.40 , $p = 0.036$; all units $10^{-4} \text{ mm}^2/\text{s}$).

As shown in Figure 2, the average FA of the cluster in the left posterior cingulum bundle showed significant negative correlations with the scores for the CPRS study problems factor ($r = -0.41$, $p = 0.019$) and the WCST total errors factor ($r = -0.41$, $p = 0.017$) and positive correlations with the WCST total correct factor ($r = 0.48$, $p = 0.005$) and the categories completed factor ($r = 0.36$, $p = 0.046$). No other significant correlation was detected.

Subgroup analysis

Our subgroup analysis of participants younger than 12 years included 26 patients (mean age 8.7 ± 1.3 yr, all boys) and

Table 1: Demographic characteristics, behavioural measures and executive functions of patients with ADHD and healthy controls

Characteristic	Group, mean \pm SD*		<i>p</i> value	Statistic
	Patients (<i>n</i> = 33)	Controls (<i>n</i> = 35)		
Age, yr	9.7 \pm 2.2	10.5 \pm 1.8	0.12	$t_{66} = -1.67$
Male:female, no.	32:1	33:2	0.59	$F_1 = 0.29$
IQ scores	104.2 \pm 5.2	107.1 \pm 6.4	0.05	$t_{66} = -1.99$
Birth weight, kg	3.3 \pm 0.8	3.6 \pm 0.9	0.26	$t_{66} = -1.14$
Parental education, yr	12.2 \pm 3.0	11.2 \pm 2.9	0.15	$t_{66} = 1.45$
Conners' Parent Rating Scale				
Conduct problem	13.1 \pm 7.4	5.6 \pm 4.9	< 0.001	$t_{54} = 4.69$
Study problem	7.7 \pm 2.9	3.1 \pm 2.3	< 0.001	$t_{66} = 6.83$
Psychosomatic	1.3 \pm 1.4	0.4 \pm 0.8	< 0.001	$t_{49} = 2.88$
Hyperactivity-impulsivity	6.2 \pm 3.0	2.4 \pm 2.0	< 0.001	$t_{54} = 5.92$
Anxiety	1.4 \pm 1.7	1.7 \pm 1.8	0.304	$t_{66} = -1.01$
Hyperactivity index	14.0 \pm 6.2	5.4 \pm 4.3	< 0.001	$t_{55} = 6.39$
Stroop Colour-Word Test				
Total time, s	291 \pm 99	186 \pm 48	< 0.001	$t_{46} = 5.40$
No. right	94.5 \pm 7.6	105.1 \pm 4.0	< 0.001	$t_{47} = -7.00$
No. errors	17.5 \pm 7.6	6.0 \pm 4.1	< 0.001	$t_{49} = 7.56$
No. corrections	9.3 \pm 5.7	3.1 \pm 2.2	< 0.001	$t_{41} = 5.79$
Wisconsin Card Sorting Test				
Total correct	27.4 \pm 10.1	33.8 \pm 5.2	0.002	$t_{47} = -3.20$
Total errors	19.2 \pm 11.7	11.0 \pm 7.2	0.001	$t_{53} = 3.42$
Perseverative errors	5.5 \pm 5.4	2.1 \pm 2.3	0.001	$t_{43} = 3.30$
Nonperseverative errors	13.8 \pm 8.0	8.9 \pm 5.6	0.006	$t_{55} = 2.81$
Categories completed	3.8 \pm 1.9	5.1 \pm 1.2	0.001	$t_{52} = -3.26$

ADHD = attention-deficit/hyperactivity disorder; SD = standard deviation.
*Unless otherwise indicated.




Fig. 1: Regions with significantly increased fractional anisotropy (FA) in patients with attention-deficit/hyperactivity disorder (ADHD). One cluster with increased FA in the left posterior cingulum bundle in patients compared with controls, corrected for multiple comparison ($p < 0.05$ after false-discovery rate correction).

23 controls (age 9.7 ± 1.2 yr, all boys); there was a significant age difference between the 2 groups ($p = 0.013$). Voxel-wise comparison of FA between the 2 groups revealed no cluster with significant FA differences after multiple comparison correction ($p < 0.05$ for FDR correction). At a relaxed threshold of $p < 0.001$ for voxel level, 1 cluster was revealed to have higher FA in patients with ADHD than controls and was located close to the cluster identified by the whole group analysis in the left posterior cingulum bundle (MNI coordinates: $x, y, z = -6, -22, 32, t = 4.17$, cluster size 20 voxels). In the whole-group analysis the larger sample size probably enhanced the effect size of this cluster so that it survived multiple comparison correction.

Our subgroup analysis of boys only included 32 patients (mean age 9.6 ± 2.3 yr) and 33 controls (mean age 10.6 ± 1.7 yr), with no significant age difference between the 2 groups ($p = 0.07$). After multiple comparison correction ($p < 0.05$ for FDR correction), 1 cluster was revealed to have elevated FA in the ADHD group and was located close to the cluster identified by the whole group analysis in the left posterior cingulum bundle (MNI coordinates: $x, y, z = -6, -22, 32, t = 5.28$, cluster size 14 voxels).

No cluster with decreased FA was found in the ADHD group in either subgroup analysis.

Discussion

Using whole brain DTI analysis we have characterized white matter microstructural abnormalities in a relatively large sample of medication-naïve children and adolescents with pure ADHD. Compared with healthy controls, patients showed increased FA in the left posterior cingulum bundle. Furthermore, the elevated FA in this cluster also showed significant correlations with behavioural measures on the CPRS and executive function scores on the WCST, supporting a role for white matter microstructure abnormalities in the pathophysiology of ADHD. This anomalous microstructure was also found in subanalyses of boys only and of children younger than 12 years, suggesting the independence of sex and maturation effects.

The cingulum bundle is the longitudinally oriented aggregate of myelinated axons underlying the cingulate cortex, whose projections connect different parts of the cingulate cortex with other brain areas, especially the medial frontal lobe, the temporal lobe and the parahippocampal gyrus.³⁴ As an important component of the limbic system, the cingulum bundle is involved in emotions and cognitive functions, including attention, memory and motivation,^{35,36} and shows abnormalities




Fig. 2: Correlations of fractional anisotropy (FA) in left posterior cingulum bundle (CB). The averaged FA of the cluster in the left posterior CB showed significant negative correlations with the study problem scores on the Conners' Parents Rating Scale (CPRS) and the total errors scores on the Wisconsin Card Sorting Test (WCST; upper panels) and positive correlations with the total correct and categories completed scores on the WCST (lower panels).

in psychiatric disorders, including schizophrenia, major depression and obsessive-compulsive disorder.^{37–39}

Previous DTI studies of ADHD have reported FA abnormalities in the cingulum bundle, but the results have been inconsistent. Four studies have reported decreased FA.^{10,40–42} Makris and colleagues¹⁰ found decreased FA in the right cingulum bundle in adults with childhood ADHD compared with healthy controls, suggesting the persistence of structural abnormalities into adulthood. Konrad and colleagues⁴⁰ replicated this finding with a larger sample of medication-naïve adult patients.⁴⁰ Two studies using diffusion spectrum imaging tractography⁴¹ and DTI tractography,⁴² respectively, also found lower FA in the bilateral cingulum bundle in youths with ADHD. In contrast, our study and 2 others^{16,32} reported increased FA of the cingulum bundle in children and adolescents with ADHD; 1 study,¹⁶ based on a detailed analysis of eigenvalues (λ_1 , λ_2 and λ_3), further proposed reduced neural branching as an explanation for this FA increase. Finally, several studies have reported no abnormalities in the cingulum bundle in patients with ADHD.^{43–45} These discrepancies might be explained by variations in methodology and demographic characteristics (e.g., medication status, different comorbidities and subtypes) of participants. To minimize such confounding, the present study included only drug-naïve patients with the combined ADHD type, and participants were carefully screened for comorbidities.

Our finding of increased FA in the cingulum bundle at the posterior cingulate gyrus region is therefore consistent with those of 2 previous studies.^{16,21} Little consideration has been given in the pathophysiology of ADHD to dysfunction of the posterior cingulum bundle, although it is a plausible hypothesis given that this carries the major efferent and afferent fibres to the posterior cingulate cortex, which participates in the regulation of attention⁴⁶ and internally directed cognition⁴⁷ and is a key component of the DMN. The DMN is a brain network associated with self-reflection, self-referential cognitions and mind-wandering. It is active in a state of rest and deactivated during tasks that make cognitive demands, with stronger deactivation corresponding to increasing attentional demands.^{48,49} Increasingly, ADHD is considered as a DMN disorder⁵⁰ in which attentional lapses derive from an inappropriate persistence of DMN activation⁴⁹; for example, a study using a stop signal task found that failure to suppress the DMN was associated with greater errors.⁵¹ Recent pathophysiological models of ADHD recognize it as a disorder encompassing large-scale neural networks, such as the cognitive control network and limbic networks, not only abnormal connectivity within the DMN itself. For instance, studies have found a negative correlation between the cognitive control network and the DMN in controls, whereas this inverse correlation was attenuated in patients with ADHD.⁵² In this conception, the interplay of the DMN with other neural networks is likely crucial to suboptimal neural functioning.⁹ As the posterior cingulum bundle is a junction containing reciprocal connections between the prefrontal cortex, anterior cingulate cortex and posterior cingulate cortex,⁵³ the abnormality we have identified is in agreement with this hypothesis.

Our correlation analysis provides further support for a pathophysiological role of the posterior cingulum bundle in patients with ADHD and is consistent with the hypothesis that components of executive functions may be associated with the limbic system in addition to the prefrontal regions.^{54,55} However, it is interesting to note that in our study patients with higher FA displayed better executive functions, albeit still worse than those of healthy controls. This phenomenon was first noted in one of our previous studies,¹⁹ in which we found a positive correlation between increased FA of right frontal white matter and the Stroop-CW Test scores in another group of children with ADHD.

As for the reason why FA is increased in patients with ADHD, further analysis of AD and RD showed that this resulted from both increased water mobility along the principal fibre direction (AD) and decreased mobility perpendicular to the fibre direction (RD), with the former component being predominant. Numerous factors, such as fibre myelination, fibre crossings, fibre density and caliber, influence AD and RD, so the exact pathological mechanism of the increased FA is uncertain. Some previous studies identifying increased FA in patients have suggested abnormally decreased dendritic branching or anomalous white matter development.^{16,21} However, as both the present study and our earlier study published in 2010¹⁹ found better executive function in patients with higher FA, we are more inclined to postulate a compensatory mechanism in children and adolescents with ADHD; for (hypothetical) example, excessive myelin hyperplasia in initially defective fibres, perhaps partially compensating for the cognitive deficits.

Limitations

Our study has some limitations. First, the sample size was small, although it was larger than those of previous whole brain DTI studies in children and adolescents with ADHD. Second, the adoption of FDR correction enhanced the impact of our result, but the small cluster size (8 voxels) limited its interpretation. Finally, a cross-sectional study cannot shed light on the development of the condition; longitudinal studies are therefore required to confirm and extend our conclusions.

Conclusion

We found that medication-naïve children and adolescents with ADHD (combined type) have increased FA in the left posterior cingulum bundle and that the FA value of this cluster was associated with behavioural problems as well as executive dysfunctions. Our results suggested that, beyond the frontal-striatal-cerebellar circuit, children and adolescents with ADHD also have pathophysiological abnormalities in the limbic network. Moreover, such microstructural alterations might be a potential substrate for abnormal behaviours and executive deficits in patients with ADHD.

Acknowledgments: This study was supported by the National Natural Science Foundation (Grant Nos. 81030027, 81227002, 81171488 and 81220108013), National Key Technologies R&D Program

(Program No. 2012BAI01B03) and Program for Changjiang Scholars and Innovative Research Team in University (Grant No. IRT1272) of China. Dr Gong would also like to acknowledge his Visiting Adjunct Professor appointment in the Department of Psychiatry at the Yale School of Medicine, Yale University, USA.

Affiliations: From the Huaxi MR Research Center, Department of Radiology, West China Hospital of Sichuan University, Chengdu, China (L. Chen, Huang, Lei, Hu, Gong); Department of Psychiatry, State Key Laboratory of Biotherapy, West China Hospital of Sichuan University, Chengdu, China (He, Y. Chen, Li, Zhou, Guo); Magnetic Resonance and Image Analysis Research Centre and Institute of Ageing and Chronic Disease, University of Liverpool, UK (Kemp).

Competing interests: None declared.

Contributors: L. Chen, X. Huang, L. Guo and Q. Gong designed the study. L. Chen, X. Huang, D. Lei, N. He, X. Hu, Y. Chen, Y. Li and J. Zhou acquired the data, which L. Chen, X. Huang, D. Lei, N. He, L. Guo, G. Kemp and Q. Gong analyzed. L. Chen and X. Huang, wrote the article, which all authors reviewed and approved for publication.

References

- Polanczyk G, de Lima MS, Horta BL, et al. The worldwide prevalence of ADHD: a systematic review and meta-regression analysis. *Am J Psychiatry* 2007;164:942-8.
- Simon V, Czobor P, Balint S, et al. Prevalence and correlates of adult attention-deficit hyperactivity disorder: meta-analysis. *Br J Psychiatry* 2009;194:204-11.
- Chamberlain SR, Robbins TW, Winder-Rhodes S, et al. Translational approaches to frontostriatal dysfunction in attention-deficit/hyperactivity disorder using a computerized neuropsychological battery. *Biol Psychiatry* 2011;69:1192-203.
- Biederman J, Faraone SV, Monuteaux MC, et al. Gender effects on attention-deficit/hyperactivity disorder in adults, revisited. *Biol Psychiatry* 2004;55:692-700.
- Nagel BJ, Bathula D, Herting M, et al. Altered white matter microstructure in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2011;50:283-92.
- Nakao T, Radua J, Rubia K, et al. Gray Matter volume abnormalities in ADHD: voxel-based meta-analysis exploring the effects of age and stimulant medication. *Am J Psychiatry* 2011;168:1154-63.
- Cortese S, Kelly C, Chabernaud C, et al. Toward systems neuroscience of ADHD: a meta-analysis of 55 fMRI studies. *Am J Psychiatry* 2012;169:1038-55.
- van Ewijk H, Heslenfeld DJ, Zwiers MP, et al. Diffusion tensor imaging in attention deficit/hyperactivity disorder: a systematic review and meta-analysis. *Neurosci Biobehav Rev* 2012;36:1093-106.
- Castellanos FX, Proal E. Large-scale brain systems in ADHD: beyond the prefrontal-striatal model. *Trends Cogn Sci* 2012;16:17-26.
- Makris N, Buka SL, Biederman J, et al. Attention and executive systems abnormalities in adults with childhood ADHD: a DT-MRI study of connections. *Cereb Cortex* 2008;18:1210-20.
- Castellanos FX, Lee PP, Sharp W, et al. Developmental trajectories of brain volume abnormalities in children and adolescents with attention-deficit/hyperactivity disorder. *JAMA* 2002;288:1740-8.
- Basser PJ. Inferring microstructural features and the physiological state of tissues from diffusion-weighted images. *NMR Biomed* 1995;8:333-44.
- Beaulieu C. The basis of anisotropic water diffusion in the nervous system — a technical review. *NMR Biomed* 2002;15:435-55.
- Giuliani NR, Calhoun VD, Pearlson GD, et al. Voxel-based morphometry versus region of interest: a comparison of two methods for analyzing gray matter differences in schizophrenia. *Schizophr Res* 2005;74:135-47.
- Ashtari M, Kumra S, Bhaskar SL, et al. Attention-deficit/hyperactivity disorder: a preliminary diffusion tensor imaging study. *Biol Psychiatry* 2005;57:448-55.
- Silk TJ, Vance A, Rinehart N, et al. White-matter abnormalities in attention deficit hyperactivity disorder: a diffusion tensor imaging study. *Hum Brain Mapp* 2009;30:2757-65.
- Davenport ND, Karatekin C, White T, et al. Differential fractional anisotropy abnormalities in adolescents with ADHD or schizophrenia. *Psychiatry Res* 2010;181:193-8.
- Kobel M, Bechtel N, Specht K, et al. Structural and functional imaging approaches in attention deficit/hyperactivity disorder: Does the temporal lobe play a key role? *Psychiatry Res* 2010;183:230-6.
- Li Q, Sun J, Guo L, et al. Increased fractional anisotropy in white matter of the right frontal region in children with attention-deficit/hyperactivity disorder: a diffusion tensor imaging study. *Neuroendocrinol Lett* 2010;31:747-53.
- Qiu MG, Ye Z, Li QY, et al. Changes of brain structure and function in ADHD children. *Brain Topogr* 2011;24:243-52.
- Peterson DJ, Ryan M, Rimrodt SL, et al. Increased regional fractional anisotropy in highly screened attention-deficit hyperactivity disorder (ADHD). *J Child Neurol* 2011;26:1296-302.
- Chuang TC, Wu MT, Huang SP, et al. Diffusion tensor imaging study of white matter fiber tracts in adolescent attention-deficit/hyperactivity disorder. *Psychiatry Res* 2013;211:186-7.
- Banaschewski T, Hollis C, Oosterlaan J, et al. Towards an understanding of unique and shared pathways in the psychopathophysiology of ADHD. *Dev Sci* 2005;8:132-40.
- Chabernaud C, Mennes M, Kelly C, et al. Dimensional brain-behavior relationships in children with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2012;71:434-42.
- Biederman J, Faraone SV. Attention-deficit hyperactivity disorder. *Lancet* 2005;366:237-48.
- Wang Z, Yang S, Jiang C, et al. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Chinese Revision. Beijing: Beijing Suicide Research and Prevention Center; 2009.
- Gong Y, Cai T. Wechsler Intelligence Scale for Children, Chinese Revision. *Chin J Clin Psychol* 1994;2:1-6.
- Annett M. A classification of hand preference by association analysis. *Br J Psychol* 1970;61:303-21.
- Conners CK, Sitarenios G, Parker JD, et al. The revised Conners' Parent Rating Scale (CPRS-R): factor structure, reliability, and criterion validity. *J Abnorm Child Psychol* 1998;26:257-68.
- Trenerry MRCB, Leber WR. Stroop Neuropsychological Screening Test. Vol 1: Psychological Assessment Resource Inc; 1989.
- Nelson HE. A modified card sorting test sensitive to frontal lobe defects. *Cortex* 1976;12:313-24.
- Van Mourik R, Oosterlaan J, Sergeant JA. The Stroop revisited: a meta-analysis of interference control in AD/HD. *J Child Psychol Psychiatry* 2005;46:150-65.
- Buchsbaum BR, Greer S, Chang WL, et al. Meta-analysis of neuroimaging studies of the Wisconsin Card-Sorting Task and component processes. *Hum Brain Mapp* 2005;25:35-45.
- Mori S, Wakana S, Van Zijl PC, et al. MRI atlas of human white matter. American Society of Neuroradiology; 2005.
- Mesulam MM. Large-scale neurocognitive networks and distributed processing for attention, language, and memory. *Ann Neurol* 1990;28:597-613.

36. Schmahmann JD, Pandya D. Fiber pathways of the brain. Oxford University Press; 2009.
37. Lochner C, Fouche JP, du Plessis S, et al. Evidence for fractional anisotropy and mean diffusivity white matter abnormalities in the internal capsule and cingulum in patients with obsessive-compulsive disorder. *J Psychiatry Neurosci* 2012;37:193-9.
38. Wang F, Sun Z, Cui L, et al. Anterior cingulum abnormalities in male patients with schizophrenia determined through diffusion tensor imaging. *Am J Psychiatry* 2004;161:573-5.
39. Zhang A, Leow A, Ajilore O, et al. Quantitative tract-specific measures of uncinate and cingulum in major depression using diffusion tensor imaging. *Neuropsychopharmacology* 2012;37:959-67.
40. Konrad A, Dielentheis TF, El Masri D, et al. Disturbed structural connectivity is related to inattention and impulsivity in adult attention deficit hyperactivity disorder. *Eur J Neurosci* 2010;31:912-9.
41. Lin HY, Gau SS, Huang-Gu SL, et al. Neural substrates of behavioral variability in attention deficit hyperactivity disorder: based on ex-Gaussian reaction time distribution and diffusion spectrum imaging tractography. *Psychol Med* 2014;44:1751-64.
42. Hong SB, Zalesky A, Fornito A, et al. Connectomic disturbances in attention-deficit/hyperactivity disorder: a whole-brain tractography analysis. *Biol Psychiatry* 2014;76:656-63.
43. Hamilton LS, Levitt JG, O'Neill J, et al. Reduced white matter integrity in attention-deficit hyperactivity disorder. *Neuroreport* 2008;19:1705-8.
44. Lawrence KE, Levitt JG, Loo SK, et al. White matter microstructure in attention-deficit/hyperactivity disorder subjects and their siblings. *J Am Acad Child Adolesc Psychiatry* 2013;52:431-40.e4.
45. Langevin LM, MacMaster FP, Crawford S, et al. Common white matter microstructure alterations in pediatric motor and attention disorders. *J Pediatrics* 2014;164(5):1157-64. e1.
46. Hahn B, Ross TJ, Stein EA. Cingulate activation increases dynamically with response speed under stimulus unpredictability. *Cereb Cortex* 2007;17:1664-71.
47. Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci* 2008;1124:1-38.
48. Hoekzema E, Carmona S, Ramos-Quiroga JA, et al. An independent components and functional connectivity analysis of resting state fMRI data points to neural network dysregulation in adult ADHD. *Hum Brain Mapp* 2014;35:1261-72.
49. Posner J, Park C, Wang Z. Connecting the dots: a review of resting connectivity MRI studies in attention-deficit/hyperactivity disorder. *Neuropsychol Rev* 2014;24:3-15.
50. Sonuga-Barke EJ, Castellanos FX. Spontaneous attentional fluctuations in impaired states and pathological conditions: a neurobiological hypothesis. *Neurosci Biobehav Rev* 2007;31:977-86.
51. Li CS, Yan P, Bergquist KL, et al. Greater activation of the "default" brain regions predicts stop signal errors. *Neuroimage* 2007;38:640-8.
52. Castellanos FX, Margulies DS, Kelly C, et al. Cingulate-precuneus interactions: a new locus of dysfunction in adult attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2008;63:332-7.
53. Jones DK, Christiansen KF, Chapman RJ, et al. Distinct subdivisions of the cingulum bundle revealed by diffusion MRI fibre tracking: implications for neuropsychological investigations. *Neuropsychologia* 2013;51:67-78.
54. Bush G, Frazier JA, Rauch SL, et al. Anterior cingulate cortex dysfunction in attention-deficit/hyperactivity disorder revealed by fMRI and the counting Stroop. *Biol Psychiatry* 1999;45:1542-52.
55. Mayberg H. Depression, II: localization of pathophysiology. *Am J Psychiatry* 2002;159:1979.

Service Information

Subscriptions and sales

Annual subscriptions are available at the following rates in 2015 (Canadian customers pay in Canadian funds and add applicable taxes; US and rest of world pay in US funds): Can\$210 or US\$247 individuals, Can\$326 or US\$378 institutions, Can\$42 for Canadian students or residents, and Can\$25 or US\$25 for single or back issues. For more information or to subscribe, please contact the Canadian Medical Association Subscription Office, PO box 830350, Birmingham AL 35283-0350; phone 800 633-4931 (Canada and Continental US only) or 205 995-1567; fax 205 995-1588; cma@subscriptionoffice.com.

Change of address

Subscribers should send the new address and the effective date of change to the CMA Member Service Centre at cmamsc@cma.ca. Allow at least 8 weeks' notice to ensure uninterrupted service.

Reprints

Commercial and author reprints can be purchased through Sheridan Press. To purchase commercial article reprints and ePrints, or to request a quote, please contact Matt Neiderer, Content Sales, Sheridan Content Services at 800 635-7181, ext. 8265; matt.neiderer@sheridan.com. Authors can order reprints by submitting an author reprint order form available at the Sheridan Press Electronic Order Centre at sheridan.com/cma/eoc or by contacting Lori Laughman, Customer Service Representative, Sheridan Reprints Services; lori.laughman@sheridan.com.

Permissions

Copyright for all material is held by 8872147 Canada Inc., wholly owned subsidiary of the Canadian Medical Association. We are a member of Access Copyright, The Canadian Copyright Licensing Agency, who may grant organizations and individuals, on our behalf, the right to respond to permissions requests. Please submit your request to Access Copyright using the online permission request service at <http://discovery.accesscopyright.ca/>. For more information on licensing or to obtain a price estimate, visit www.accesscopyright.ca/permissions/

Microform, abstracting and indexing

The journal is abstracted or indexed in: MEDLINE/Index Medicus, Current Contents, Biosearch Index, Social Sciences Citation Index, Biological Abstracts, EMBASE/Excerpta Medica, e-psyche, Mental Health Abstracts, ISI Web of Science, Child Development Abstracts and Bibliography, Standard Periodical Directory, Arts and Humanities Citation Index, Research Alert (formerly Automated Subject Citation Alert), Criminal Justice Abstracts, Neuroscience Citation Index, Psychological Abstracts, Sociological Abstracts. *JPN* is available on microform from ProQuest Information and Learning, PO Box 1346, Ann Arbor MI 48106-1346; il.proquest.com.