- Murray E, Brouwer S, McCutcheon R, et al. Opposing neural effects of naltrexone on food reward and aversion: implications for the treatment of obesity. *Psychopharmacology (Berl)* 2014;231:4323-35.
- Koob GF, Rocio M, Carrera A, et al. Substance dependence as a compulsive behavior. J Psychopharmacol 1998;12:39-48.
- Hutcheson DM, Everitt BJ, Robbins TW, et al. The role of withdrawal in heroin addiction: enhances reward or promotes avoidance? *Nat Neurosci* 2001;4:943-7.
- Grace AA. The tonic/phasic model of dopamine system regulation: its relevance for understanding how stimulant abuse can alter basal ganglia function. *Drug Alcohol Depend* 1995;37:111-29.
- Kelley AE. Ventral striatal control of appetitive motivation: role in ingestive behavior and reward-related learning. *Neurosci Biobehav Rev* 2004;27:765-76.
- Kringelbach ML, Rolls ET. The functional neuroanatomy of the human orbitofrontal cortex: evidence from neuroimaging and neuropsychology. *Prog Neurobiol* 2004;72:341-72.
- Schoenbaum G, Roesch MR, Stalnaker TA. Orbitofrontal cortex, decision-making and drug addiction. *Trends Neurosci* 2006;29:116-24.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Text Revision (DSM-IV-TR). Washington, DC: APA; 2000.
- 33. McLellan AT, Kushner H, Metzger D, et al. The fifth edition of the Addiction Severity Index. J Subst Abuse Treat 1992;9:199-213.
- Wesson DR, Ling W. The clinical opiate withdrawal scale (COWS). J Psychoactive Drugs 2003;35:253-9.
- Filbey FM, Claus E, Audette AR, et al. Exposure to the taste of alcohol elicits activation of the mesocorticolimbic neurocircuitry. *Neuropsychopharmacology* 2008;33:1391-401.
- 36. Filbey FM, Schacht JP, Myers US, et al. Marijuana craving in the brain. *Proc Natl Acad Sci U S A* 2009;106:13016-21.
- Volkow ND, Wang GJ, Telang F, et al. Cocaine cues and dopamine in dorsal striatum: mechanism of craving in cocaine addiction. J Neurosci 2006;26:6583-8.
- Crockford DN, Goodyear B, Edwards J, et al. Cue-induced brain activity in pathological gamblers. *Biol Psychiatry* 2005;58:787-95.
- Krupitsky E, Nunes EV, Ling W, et al. Injectable extended-release naltrexone for opioid dependence: a double-blind, placebo-controlled, multicentre randomised trial. *Lancet* 2011;377:1506-13.
- 40. O'Brien CP, Gastfriend DR, Forman RF, et al. Long-term opioid blockade and hedonic response: preliminary data from two openlabel extension studies with extended-release naltrexone. *Am J Addict* 2011;20:106-12.
- Lieberman MD, Cunningham WA. Type I and Type II error concerns in fMRI research: re-balancing the scale. Soc Cogn Affect Neurosci 2009;4:423-8.
- Nichols TE, Eklund A, Knutsson H. A defense of using restingstate fMRI as null data for estimating false positive rates. *Cogn Neurosci* 2017;8:144-9.

- Slotnick SD. Cluster success: fMRI inferences for spatial extent have acceptable false-positive rates. Cogn Neurosci 2017;8:150-5.
- 44. Slotnick SD. Resting-state fMRI data reflects default network activity rather than null data: a defense of commonly employed methods to correct for multiple comparisons. *Cogn Neurosci* 2017;8:141-3.
- Eklund A, Nichols TE, Knutsson H. Cluster failure: why fMRI inferences for spatial extent have inflated false-positive rates. *Proc Natl Acad Sci U S A* 2016;113:7900-5.
- Boecker H, Sprenger T, Spilker ME, et al. The runner's high: opioidergic mechanisms in the human brain. *Cereb Cortex* 2008; 18:2523-31.
- Navratilova E, Atcherley CW, Porreca F. Brain circuits encoding reward from pain relief. *Trends Neurosci* 2015;38:741-50.
- Robinson TE, Berridge KC. The psychology and neurobiology of addiction: an incentive-sensitization view. *Addiction* 2000;95(Suppl 2):S91-117.
- Everitt BJ, Robbins TW. Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci* 2005;8:1481-9.
- Lukas SE, Lowen SB, Lindsey KP, et al. Extended-release naltrexone (XR-NTX) attenuates brain responses to alcohol cues in alcohol-dependent volunteers: a bold FMRI study. *Neuroimage* 2013;78:176-85.
- Meller RE, Keverne EB, Herbert J. Behavioural and endocrine effects of naltrexone in male talapoin monkeys. *Pharmacol Biochem Behav* 1980;13:663-72.
- Shayit M, Nowak R, Keller M, et al. Establishment of a preference by the newborn lamb for its mother: the role of opioids. *Behav Neurosci* 2003;117:446-54.
- Inagaki TK, Ray LA, Irwin MR, et al. Opioids and social bonding: naltrexone reduces feelings of social connection. *Soc Cogn Affect Neurosci* 2016;11:728-35.
- Wang AL, Lowen SB, Elman I, et al. Sustained opioid antagonism increases striatal sensitivity to baby schema in opioid dependent women. *Eur Neuropsychopharmacol* 2016;26:S695-6.
- 55. United Nations Office on Drugs and Crime. *World Drug Report* 2014. Vienna, Austria: United Nations; 2014.
- Cicero TJ, Ellis MS, Surratt HL, et al. The changing face of heroin use in the United States: a retrospective analysis of the past 50 years. JAMA Psychiatry 2014;71:821-6.
- Sullivan MA, Bisaga A, Mariani JJ, et al. Naltrexone treatment for opioid dependence: Does its effectiveness depend on testing the blockade? *Drug Alcohol Depend* 2013;133:80-5.
- Bradley MM, Lang PJ. Measuring emotion: the self-assessment manikin and the semantic differential. J Behav Ther Exp Psychiatry 1994;25:49-59.
- Kim BW, Kennedy DN, Lehar J, et al. Recurrent, robust and scalable patterns underlie human approach and avoidance. *PLoS ONE* 2010;5:e10613.

Correction: Enhanced corticobulbar excitability in chronic smokers during visual exposure to cigarette smoking cues

In the article "Enhanced corticobulbar excitability in chronic smokers during visual exposure to cigarette smoking cues" by Vicario and colleagues¹ published in the July 2014 issue of the journal, there was an error in the legend of Figure 1. The dark grey and light grey bars were labelled as the smoking cue and scramble, respectively; however, the light grey bars should have been labelled as the smoking cue and the dark grey bars as the scramble. The figure has been updated in the online version of the article, which is available at jpn.ca.

We apologize for the error.

Reference

1. Vicario CM, Komeilipoor N, Cesari P, et al. Enhanced corticobulbar excitability in chronic smokers during visual exposure to cigarette smoking cues. *J Psychiatry Neurosci* 2014;39:232–8.