

**Appendix 1** to Miskowiak KW, Forman J, Vinberg M, et al. Impact of pre-treatment inter-hemispheric hippocampal asymmetry on improvement in verbal learning following erythropoietin treatment in mood disorders: a randomized controlled trial. *J Psychiatry Neurosci* 2019.

DOI: 10.1503/jpn.180205

Copyright © 2019 The Author(s) or their employer(s). To receive this resource in an accessible format, please contact us at [cmajgroup@cmaj.ca](mailto:cmajgroup@cmaj.ca).

*Online appendices are unedited and posted as supplied by the authors.*

**Supplementary figure 1.** Statistical map of relative surface area. The linear regression slope for changes in **verbal learning** was significantly different for the EPO and saline groups within the PFC ROI ( $p < 0.02$ ). Specifically, larger precentral surface area (marked with yellow and orange) was associated with greater **verbal learning** improvement in EPO- vs. saline-treated participants. The lower panels show partial regression plots for baseline to follow-up change in RAVLT scores adjusted for age sex and TIV.

**Appendix 1** to Miskowiak KW, Forman J, Vinberg M, et al. Impact of pre-treatment inter-hemispheric hippocampal asymmetry on improvement in verbal learning following erythropoietin treatment in mood disorders: a randomized controlled trial. *J Psychiatry Neurosci* 2019.

DOI: 10.1503/jpn.180205

Online appendices are unedited and posted as supplied by the authors.

