DOI: 10.1503/jpn.160219

Copyright © 2017 The Author(s) or their employer(s). To receive this resource in an accessible format, please contact us at cmajgroup@cmaj.ca.

Online appendices are unedited and posted as supplied by the authors.

Table S1

Imaging Methodology Quality Assessment Checklist

Category 1: Subjects

- 1. Patients were evaluated prospectively, specific diagnostic criteria were applied, and demographic data was reported
- 2. Healthy comparison subjects were evaluated prospectively, psychiatric and medical illnesses were excluded, and demographic data was reported
- 3. Important variables (e.g. illness duration, severity of illness, drug status, non-drug therapy status) were checked either by stratification or statistically
- 4. Sample size per group > 10, and no significant difference in age and sex existed

Category 2: Methods for image acquisition and analysis

- 5. Magnet strength at least 1.5T
- 6. Whole brain analysis was automated with no a-priori regional selection

| Appendix 1 to Gao X, Zhang W, Yao L, et al. Association between structural and functional brain alterations in drug-free patients with schizophrenia: a multimodal meta-analysis. <i>J Psychiatry Neurosci</i> 2017. DOI: 10.1503/jpn.160219 |
|---|
| Online appendices are unedited and posted as supplied by the authors. |
| 7. Coordinates reported in a standard space |
| 8. The imaging technique used was clearly described so as it could be reproduced |
| 9. Measurements were clearly described so that they could be reproduced |
| 10. Results have been corrected for multiple comparison |
| Category 3: Results and conclusions |
| 11. Statistical parameters for significant and important non-significant differences were provided |
| 12. Conclusions were consistent with the results obtained and the limitations were discussed |

DOI: 10.1503/jpn.160219

Table S2Meta-analysis results of structural MRI

| | MNI | SDM-Z | P | Voxels | Cluster breakdown (number of voxels) |
|---|---|--------|-------------|--------|--|
| Cluster of decreased gray matter volume (patien | nts <controls)< td=""><td></td><td></td><td></td><td></td></controls)<> | | | | |
| Left fusiform gyrus | -26,-54,-16 | -2.584 | 0.000154853 | 771 | Left fusiform gyrus (382) |
| | | | | | Left cerebellum, hemispheric lobule VI (182) |
| | | | | | Left parahippocampal gyrus (21) |
| | | | | | Left hippocampus (15) |
| Left inferior frontal gyrus | -46,16,0 | -2.497 | 0.00023222 | 649 | Left inferior frontal gyrus (314) |
| | | | | | Left insula (166) |

DOI: 10.1503/jpn.160219

| | | | | | Left temporal pole, superior temporal gyrus (106) |
|--|------------|--------|-------------|-----|---|
| Right superior temporal gyrus | 52,-6,-6 | -2.098 | 0.002043664 | 330 | Right superior temporal gyrus (124) |
| | | | | | Right insula (97) |
| | | | | | Right rolandic operculum (12) |
| | | | | | Right middle temporal gyrus (11) |
| | | | | | Right temporal pole, superior temporal gyrus (10) |
| Left supramarginal gyrus | -56,-40,36 | -2.387 | 0.000490248 | 293 | Left supramarginal gyrus (117) |
| | | | | | Left inferior parietal gyrus (104) |
| | | | | | Left superior temporal gyrus (64) |
| Cluster of increased gray matter volume (patients>co | ntrols) | | | | |
| Right lingual gyrus | 6,-34,-6 | 1.895 | 0.000005186 | 466 | Right lingual gyrus (62) |
| | | | | | Cerebellum, vermic lobule III (36) |

DOI: 10.1503/jpn.160219

Online appendices are unedited and posted as supplied by the authors.

Left cerebellum, hemispheric lobule IV/V (31)

Left lingual gyrus (18)

Right superior frontal gyrus, orbital part 6,60,-22 1.617 0.000805080 155 Right gyrus rectus (79)

Right superior frontal gyrus (38)

(a) Voxel probability threshold: *P*=0.005

(b) Peak height threshold: z=1

(c) Cluster extent threshold: 100 voxels. Regions with less than 10 voxels are not reported in the cluster breakdown.

DOI: 10.1503/jpn.160219

Table S3Analyses of subgroups and sensitivity analyses of structural studies

| | Increased g | ray matter volume | Decreased gray matter volume | | | |
|--|---------------|------------------------|------------------------------|---------------|----------------|--------------------|
| Subgroup analysis | Right lingual | Right superior frontal | Left fusiform | Left inferior | Right superior | Left supramarginal |
| | gyrus | gyrus | gyrus | frontal gyrus | temporal gyrus | gyrus |
| Studies with slice thickness ≤ 1.5 mm at acquisition (n=13) | YES | YES | NO | NO | YES | YES |
| Studies using $\leq 8 \text{ mm smoothing kernel (n=10)}$ | YES | YES | YES | NO | NO | YES |
| Studies using 1.5T MRI (n=10) | NO | NO | YES | YES | YES | NO |
| Studies with an additional correction step (n=9) | YES | YES | YES | YES | YES | YES |
| Studies of drug-naive patients (n=14) | YES | YES | YES | YES | YES | YES |

DOI: 10.1503/jpn.160219

Online appendices are unedited and posted as supplied by the authors.

Jackknife sensitivity analysis, discarded study

| Jackkinie sensitivity analysis, discarded study | | | | | | |
|---|-----|-----|-----|-----|-----|-----|
| 2003Salgado-Pineda | YES | YES | YES | YES | YES | YES |
| 2005Jayakumar | YES | YES | YES | YES | YES | YES |
| 2007Prasad | YES | YES | YES | YES | YES | YES |
| 2007Chua | YES | YES | YES | YES | YES | YES |
| 2008Meda | YES | YES | YES | YES | YES | YES |
| 2008Witthaus | YES | YES | YES | YES | YES | YES |
| 2010Venkatasubramanian | YES | YES | YES | YES | YES | YES |
| 2011Berge | YES | YES | YES | YES | YES | YES |
| 2013Suazo | YES | YES | YES | YES | YES | YES |
| 2013WentingRen | YES | YES | YES | YES | YES | YES |

DOI: 10.1503/jpn.160219

| 2013XiaofengGuo(S) | YES | YES | YES | YES | YES | YES |
|--------------------|-----|-----|-----|-----|-----|-----|
| 2013XiaofengGuo(L) | YES | YES | YES | YES | YES | YES |
| 2014WenbinGuo | YES | YES | YES | YES | YES | YES |
| 2014JunLi | YES | YES | YES | YES | YES | YES |
| 2014Nenadic | YES | YES | YES | YES | YES | YES |
| 2016Yue | YES | YES | YES | YES | YES | YES |

DOI: 10.1503/jpn.160219

Table S4Meta-analysis results of functional MRI

| | MNI | SDM-Z | P | Voxels | Cluster breakdown (number of voxels) |
|--|-----------|--------|-------------|--------|---|
| Cluster of hypoactivation (patients <controls)< th=""><th></th><th></th><th></th><th></th><th></th></controls)<> | | | | | |
| Right inferior frontal gyrus, opercular part | 46,10,30 | -3.221 | 0.000006437 | 893 | Right inferior frontal gyrus, opercular part (432) |
| | | | | | Right inferior frontal gyrus, triangular part (152) |
| | | | | | Right precentral gyrus (98) |
| | | | | | Right middle frontal gyrus (17) |
| Right angular gyrus | 40,-60,52 | -2.719 | 0.000062704 | 331 | Right angular gyrus (166) |
| | | | | | Right inferior parietal (excluding supramarginal and angular) gyrus (140) |

DOI: 10.1503/jpn.160219

| Unline appendices are unedited | and posted as supplied | i by the au | lnors. | | |
|---------------------------------------|------------------------|-------------|-------------|------|---|
| | | | | | Right superior parietal gyrus (25) |
| Right insula | 38,14,-14 | -2.439 | 0.000266016 | 339 | Right insula (197) |
| | | | | | Right inferior frontal gyrus, orbital part (44) |
| | | | | | Right temporal pole, superior temporal gyrus (30) |
| Cluster of hyperactivation (patients: | >controls) | | | | |
| Left striatum | -28,-2,4 | 3.273 | ~0 | 2761 | Left superior temporal gyrus (685) |
| | | | | | Left insula (564) |
| | | | | | Left rolandic operculum (335) |
| | | | | | Left lenticular nucleus, putamen (296) |
| | | | | | Left striatum (189) |
| | | | | | Left heschl gyrus (164) |
| | | | | | Left postcentral gyrus (69) |

DOI: 10.1503/jpn.160219

Online appendices are unedited and posted as supplied by the authors.

| | | | | | | Left supramarginal gyrus (44) |
|---|------------------------------|-------------|-------|-------------|-----|--|
| | | | | | | Left pons (17) |
| | Right striatum | 26,-2,8 | 1.919 | 0.000929713 | 212 | Right insula (71) |
| | | | | | | Right striatum (48) |
| | | | | | | Right lenticular nucleus, putamen (34) |
| L | eft inferior occipital gyrus | -34,-76,-12 | 2.178 | 0.000299335 | 168 | Left fusiform gyrus (83) |
| | | | | | | Left inferior occipital gyrus (42) |

(a) Voxel probability threshold: P=0.005

(b) Peak height threshold: z=1

(c) Cluster extent threshold: 100 voxels. Regions with less than 10 voxels are not reported in the cluster breakdown.

DOI: 10.1503/jpn.160219

Table S5Analyses of subgroups and sensitivity analyses of functional studies

| | | Hyperactivation | | Hypoactivation | | | |
|---|---------------|-----------------|--|-----------------------|---------------|--------|--|
| Subgroup analysis | Left striatum | Right striatum | Left inferior Right inferior frontal Right angul | | Right angular | Right | |
| | | | occipital gyrus | gyrus, opercular part | gyrus | insula | |
| Studies with slice thickness <5 mm at acquisition (n=13) | YES | NO | NO | YES | NO | YES | |
| Studies using $\leq 8 \text{ mm}$ smoothing kernel (n=15) | YES | YES | YES | YES | YES | YES | |
| Studies using 1.5T MRI (n=8) | YES | NO | NO | YES | NO | NO | |
| Studies with an additional correction step (n=16) | YES | YES | YES | YES | YES | YES | |
| Studies performing task (n=11) | YES | NO | NO | YES | NO | YES | |

DOI: 10.1503/jpn.160219

| Studies of drug-naive patients (n=8) | YES | YES | YES | YES | YES | NO |
|---|-----|-----|-----|-----|-----|-----|
| Jackknife sensitivity analysis, discarded study | | | | | | |
| 1997Andreason | YES | YES | YES | YES | YES | YES |
| 2010Scheef | YES | YES | YES | YES | YES | YES |
| 2013Ren | YES | NO | NO | YES | NO | YES |
| 2015Guo | YES | YES | YES | YES | YES | YES |
| 2014 Hadley | YES | YES | YES | YES | YES | YES |
| 2003Hofer | YES | YES | YES | YES | YES | YES |
| 2004Jones | YES | YES | YES | YES | YES | YES |
| 2005Weiss | YES | YES | YES | YES | YES | YES |
| 2005Boksman | YES | YES | YES | YES | YES | YES |

DOI: 10.1503/jpn.160219

| 2008Scheuerecker | YES | YES | YES | YES | YES | YES |
|--------------------------|-----|-----|-----|-----|-----|-----|
| 2010Comilo | YES | YES | YES | YES | YES | YES |
| 2011Nejad | YES | YES | YES | YES | YES | YES |
| 2013Bin | YES | YES | YES | YES | YES | YES |
| 2014Schlagenhauf | YES | YES | YES | YES | YES | YES |
| 2015Lesh | YES | YES | YES | YES | YES | YES |
| 2016Zheng | YES | YES | YES | YES | YES | YES |
| 2016Cui (1) ⁱ | YES | YES | YES | YES | YES | YES |
| 2016Cui (2) | YES | YES | YES | YES | YES | YES |

DOI: 10.1503/jpn.160219

Table S7

Table S6

Regions with altered gray matter volume showing significant statistical heterogeneity between studies (voxelwise p<0.005)


| Description | MNI coordinate | SDM-Z | P | Voxels |
|---------------------|----------------|-------|---------|--------|
| Right insula | 38,-16,-2 | 4.039 | ~0 | 1372 |
| Left fusiform gyrus | -36,-60,-16 | 2.414 | 0.00026 | 264 |

Regions with altered regional metabolism showing significant statistical heterogeneity between studies (voxelwise p<0.005)


| Description | MNI coordinate | SDM-Z | P | Voxels |
|-------------|----------------|-------|---------|--------|
| Left insula | -36,-6,10 | 2.209 | 0.00155 | 184 |

DOI: 10.1503/jpn.160219


Table S8Results of the funnel plots and Eager test




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219




DOI: 10.1503/jpn.160219



DOI: 10.1503/jpn.160219



DOI: 10.1503/jpn.160219



DOI: 10.1503/jpn.160219

Online appendices are unedited and posted as supplied by the authors.

References of the inclusion studies¹⁻²⁹:

- 1. Salgado-Pineda P, Baeza I, Pérez-Gómez M, et al. Sustained attention impairment correlates to gray matter decreases in first episode neuroleptic-naive schizophrenic patients[J]. *NeuroImage* 2003,19(2):365-375.
- 2. Jayakumar PN, Venkatasubramanian G, Gangadhar BN, et al. Optimized voxel-based morphometry of gray matter volume in first-episode, antipsychotic-naive schizophrenia[J]. *Progress in neuro-psychopharmacology & biological psychiatry* 2005,29(4):587-591.
- 3. Chua SE, Cheung C, Cheung V, et al. Cerebral grey, white matter and csf in never-medicated, first-episode schizophrenia[J]. *Schizophrenia research* 2007,89(1-3):12-21.
- 4. Prasad KM, Shirts BH, Yolken RH, et al. Brain morphological changes associated with exposure to HSV1 in first-episode schizophrenia[J]. *Molecular psychiatry* 2007,12(1):105-113, 101.
- 5. Meda SA, Giuliani NR, Calhoun VD, et al. A large scale (N=400) investigation of gray matter differences in schizophrenia using optimized voxel-based morphometry[J]. *Schizophrenia research* 2008,101(1-3):95-105.
- 6. Witthaus H, Kaufmann C Fau Bohner G, Bohner G Fau Ozgurdal S, et al. Gray matter abnormalities in subjects at ultra-high risk for schizophrenia and first-episode schizophrenic patients compared to healthy controls[J]. (0165-1781 (Print)).
- 7. Venkatasubramanian G. Neuroanatomical correlates of psychopathology in antipsychotic-naive schizophrenia[J]. *Indian journal of psychiatry* 2010,52(1):28-36.
- 8. Berge D, Carmona S, Rovira M, et al. Gray matter volume deficits and correlation with insight and negative symptoms in first-psychotic-episode subjects[J]. *Acta psychiatrica Scandinavica* 2011,123(6):431-439.
- 9. Suazo V, Diez A, Montes C, et al. Structural correlates of cognitive deficit and elevated gamma noise power in schizophrenia[J]. *Psychiatry and clinical neurosciences* 2014,68(3):206-215.
- 10. Ren W, Lui S, Deng W, et al. Anatomical and functional brain abnormalities in drug-naive first-episode schizophrenia[J]. *The American journal of psychiatry* 2013,170(11):1308-1316.
- 11. Guo X, Li J, Wei Q, et al. Duration of untreated psychosis is associated with temporal and occipitotemporal gray matter volume decrease in treatment naive


DOI: 10.1503/jpn.160219

- schizophrenia[J]. PloS one 2013,8(12):e83679.
- 12. Guo W, Liu F, Xiao C, et al. Dissociation of anatomical and functional alterations of the default-mode network in first-episode, drug-naive schizophrenia[J]. *Clin Neurophysiol* 2015.
- 13. Guo X, Li J, Wang J, et al. Hippocampal and orbital inferior frontal gray matter volume abnormalities and cognitive deficit in treatment-naive, first-episode patients with schizophrenia[J]. *Schizophrenia research* 2014,152(2-3):339-343.
- 14. Nenadic I, Dietzek M, Schonfeld N, et al. Brain structure in people at ultra-high risk of psychosis, patients with first-episode schizophrenia, and healthy controls: a VBM study[J]. *Schizophrenia research* 2015,161(2-3):169-176.
- 15. Scheef L, Manka C, Daamen M, et al. Resting-state perfusion in nonmedicated schizophrenic patients: a continuous arterial spin-labeling 3.0-T MR study[J]. Radiology 2010,256(1):253-260.
- 16. Hadley JA, Nenert R, Kraguljac NV, et al. Ventral tegmental area/midbrain functional connectivity and response to antipsychotic medication in schizophrenia[J]. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* 2014,39(4):1020-1030.
- 17. Hofer A, Weiss EM, Golaszewski SM, et al. Neural correlates of episodic encoding and recognition of words in unmedicated patients during an acute episode of schizophrenia: a functional MRI study[J]. *The American journal of psychiatry* 2003,160(10):1802-1808.
- 18. Jones HM, Brammer MJ, O'Toole M, et al. Cortical effects of quetiapine in first-episode schizophrenia: a preliminary functional magnetic resonance imaging study[J]. Biological psychiatry 2004,56(12):938-942.
- 19. Boksman K, Theberge J, Williamson P, et al. A 4.0-T fMRI study of brain connectivity during word fluency in first-episode schizophrenia[J]. *Schizophrenia research* 2005,75(2-3):247-263.
- 20. Weiss EM, Siedentopf C, Golaszewski S, et al. Brain activation patterns during a selective attention test--a functional MRI study in healthy volunteers and unmedicated patients during an acute episode of schizophrenia[J]. *Psychiatry research* 2007,154(1):31-40.
- 21. Scheuerecker J, Ufer S, Zipse M, et al. Cerebral changes and cognitive dysfunctions in medication-free schizophrenia an fMRI study[J]. *Journal of psychiatric research* 2008,42(6):469-476.
- 22. De la Fuente-Sandoval C, Favila R, Gomez-Martin D, et al. Functional magnetic resonance imaging response to experimental pain in drug-free patients with schizophrenia[J]. *Psychiatry research* 2010,183(2):99-104.
- 23. Nejad AB, Ebdrup BH, Siebner HR, et al. Impaired temporoparietal deactivation with working memory load in antipsychotic-naive patients with first-episode

DOI: 10.1503/jpn.160219

- schizophrenia[J]. The world journal of biological psychiatry: the official journal of the World Federation of Societies of Biological Psychiatry 2011,12(4):271-281.
- 24. Ji B, Mei W, Zhang JX, et al. Abnormal auditory sensory gating-out in first-episode and never-medicated paranoid schizophrenia patients: an fMRI study[J]. Experimental brain research 2013,229(2):139-147.
- 25. Schlagenhauf F, Huys QJ, Deserno L, et al. Striatal dysfunction during reversal learning in unmedicated schizophrenia patients[J]. *Neuroimage* 2014,89:171-180.
- 26. Lesh TA, Tanase C, Geib BR, et al. A multimodal analysis of antipsychotic effects on brain structure and function in first-episode schizophrenia[J]. *JAMA psychiatry* 2015,72(3):226-234.
- 27. Yue Y, Kong L, Wang J, et al. Regional Abnormality of Grey Matter in Schizophrenia: Effect from the Illness or Treatment?[J]. *PloS one* 2016,11(1):e0147204.
- 28. Cui LB, Liu K, Li C, et al. Putamen-related regional and network functional deficits in first-episode schizophrenia with auditory verbal hallucinations[J]. *Schizophrenia research* 2016,173(1-2):13-22.
- 29. Zheng J, Zhang Y, Guo X, et al. Disrupted amplitude of low-frequency fluctuations in antipsychotic-naive adolescents with early-onset schizophrenia[J]. *Psychiatry research* 2016,249:20-26.

DOI: 10.1503/jpn.160219



DOI: 10.1503/jpn.160219

